

Drugs Used to Treat or Control Coccidiosis in Livestock and Poultry

Not all drugs listed are approved for use in sheep and goats. The mention of non-approved products is for informational purposes only. They require extra label use and can only be used legally under the advice of a licensed veterinarian in the context of a valid veterinary-patient-client relationship.

Trade Name	Drug Ingredient	Approved Species	Specificity	Labeled Dosage	Withdrawal Time
For treatment of coccidiosis					
Albon® 12.5% solution	Sulfadimethoxine	Beef Non-lactating dairy cattle Poultry	shipping fever complex, bacterial pneumonia, calf diphtheria, foot rot, and coccidiosis (poultry)	1.18-2.36 g/gallon	Cattle: 7 days slaughter Poultry: 5 days slaughter
Corid 20% Soluble Powder	Amprolium	Cattle Chickens Turkeys	Treatment of Coccidiosis	4 oz/50 gal water (treatment) 4 oz/100 gal water (prevention)	Cattle: 24 hours
Corid 9.6% solution Amprovine	Amprolium	Cattle Poultry	For the control and treatment of outbreaks of Coccidiosis	5 mg/1 kg	Cattle: 1 day slaughter Poultry: 0 days slaughter
Di-Methox	Sulfadimethoxine	Cattle Poultry	shipping fever complex, bacterial pneumonia, calf diphtheria, and foot rot and coccidiosis (poultry)	1.18-2.36 g/gallon 2.5 g/100 lbs. day 1 1.25 g/100 lbs. day 2-5	Cattle: 7 days slaughter Chicken and Turkey: 5 days slaughter
20% Sulfaquinoxoline	Sulfaquinoxoline Sodium	Cattle Poultry	For the control and treatment of outbreaks of coccidiosis	1 oz./10 gallon water	Non-lactating cattle: 10 days Turkey and Chicken: 10 days
Sulmet® oblet	Sulfamethazine	Beef Non-lactating dairy cattle Horses	Shipping fever, bacterial scours, foot rot, calf diphtheria, acute mastitis, acute metritis and coccidiosis	100 mg /lb body weight	Cattle: 10 days
Sulmet®12.5% solution	Sulfamethazine	Cattle Swine Poultry	shipping fever complex, bacterial scours, foot rot, calf diphtheria, acute mastitis metritis and coccidiosis (poultry)	112.5 mg/lb - day 1. 56.25/mg/lb	Cattle: 10 days slaughter Swine: 15 days slaughter Poultry: 10 days

For prevention of coccidiosis					
Bovatec® Premix	Lasalocid Sodium	Beef Dairy heifers Poultry Sheep in confinement Rabbits TOXIC TO EQUINES	<i>Eimeria ovina</i> <i>E. crandallis</i> <i>E. ovinoidalis</i> (<i>E. ninakohlyakimovae</i>) <i>E. parva</i> , and <i>E. intricata</i>	20-30 g/ton feed	All species: 0 days slaughter
Deccox®	Decoquinatate	Beef non-lactating dairy cattle Chickens Non-lactating goats Non-lactating lambs	<i>E. ovinoidalis</i> , <i>E. crandallis</i> , <i>E. parva</i> , <i>E. bakuensis</i> , <i>E. christensenii</i> , and <i>E. ninakohlyakimovae</i>	13.6 g/ton feed	All species: 0 days slaughter
Rumensin®	Monensin Sodium	Beef Dairy heifers Non-lactating goats TOXIC TO EQUINES	<i>Eimeria crandallis</i> <i>E. christensenii</i> <i>E. ninakohlyakimovae</i>	20 g/ton feed	cattle: 0 days slaughter goats: 0 days slaughter

Source: Product labels and FDA Approved Animal Drug Products Online Database System @ <http://dil.vetmed.vt.edu/>.

Last Updated 23-Sep-2004 by Susan Schoenian, Area Agent, Sheep & Goats, Western Maryland Research & Education Center, Maryland Cooperative Extension.

Copyright. 2004. Maryland Small Ruminant Page.



<http://www.agnr.umd.edu/mce>